arXiv:1603.05518v1 [math.OA] 16 Mar 2016

ON CONNECTED LIE GROUPS AND THE APPROXIMATION PROPERTY

SØREN KNUDBY

ABSTRACT. Recently, a complete characterization of connected Lie groups with the Approximation Property was given. The proof used the newly introduced property (T*). We present here a short proof of the same result avoiding the use of property (T*). Using property (T*), however, the characterization is extended to every almost connected group. We end with some remarks about the impossibility of going beyond the almost connected case.

The Fourier algebra $A(G)$ of a locally compact group G was introduced by Eymard [5] and can be defined as the coeffient space of the left regular representation of G on $L^2(G)$,

$$A(G) = \{f * \widetilde{g} \mid f, g \in L^2(G)\},$$

where $\widetilde{g}(x) = \overline{g(x^{-1})}$. One can identify $A(G)$ with the predual of the group von Neumann algebra associated with the left regular representation. A multiplier φ of $A(G)$ is called completely bounded if its transposed operator M_φ is a completely bounded operator on the group von Neumann algebra. The completely bounded multiplier norm of φ is defined as

$$\|\varphi\|_{M_0A} = \|M_\varphi\|_{\mathrm{cb}}.$$

The space of completely bounded multipliers is dentoed $M_0A(G)$. One has an inclusion $A(G) \subseteq M_0A(G)$. It was shown in [4, Proposition 1.10] that the space $M_0A(G)$ has a predual obtained as the completion of $L^1(G)$ in the norm

$$\|f\|_{M_0A(G)_*} = \sup\left\{\left|\int_G f(x)\varphi(x)\,dx\right| : \varphi \in M_0A(G),\ \|\varphi\|_{M_0A} \leq 1\right\}.$$

Remark 1. In general, $M_0A(G)$ does not have a unique predual nor a unique weak* topology. The predual always refers to the one constructed explicitly above, and the weak* topology on $M_0A(G)$ is the weak* topology coming from this explicit predual.

Following Haagerup and Kraus [9], a locally compact group G has the Approximation Property (short: AP) if there exists a net in $A(G)$ converging to 1 in the weak* topology of $M_0A(G)$. Haagerup and Kraus showed that many groups have this property, e.g., all weakly amenable groups. This includes solvable groups, compact groups, and simple Lie groups of real rank one [3, 10]. They showed moreover that the AP is preserved by passing to closed subgroups and preserved under group extensions (as opposed to weak amenability). It is also well-known (and routine to verify) that if K is a compact normal subgroup in G, then G has the AP if and only if G/K has the AP. We refer to [3, 4, 9] for details on completely bounded multipliers and the AP.

Date: March 18, 2016.
Supported by the Deutsche Forschungsgemeinschaft through the Collaborative Research Centre (SFB 878).

Haagerup and Kraus conjectured that the group $\mathrm{SL}(3,\mathbb{R})$ does not have the AP, but the conjecture was settled only much later by Lafforgue and de la Salle [11]. Not long after, Haagerup and de Laat showed more generally that any simple Lie group of real rank at least two does not have the AP [6, 7]. Finally, a complete characterization of connected Lie groups with the AP was given in [8] (see also Theorem 2 below). The proof in [8] involved among other things the property (T*), introduced in [8]. We give below a much shorter proof of the characterization of connected Lie groups with the AP that avoids the use of property (T*).

To state the characterization of connected Lie groups with the AP, we first recall the Levi decomposition of such a group (see [15, Section 3.18] for details). For a connected Lie group G, one can decompose its Lie algebra $\mathfrak{g}$ as $\mathfrak{g} = \mathfrak{r} \rtimes \mathfrak{s}$, where $\mathfrak{r}$ is the solvable radical and $\mathfrak{s}$ is a semisimple subalgebra. One can further decompose $\mathfrak{s} = \mathfrak{s}_1 \oplus \cdots \oplus \mathfrak{s}_n$ into simple summands $\mathfrak{s}_i$ ($i = 1, \ldots, n$). Let R, S, and S_i denote the corresponding connected Lie subgroups of G. One has $G = RS$ as a set. This is called a Levi decomposition of G.

The subgroup R is solvable, normal, and closed. The subgroup S is semisimple and locally isomorphic to the direct product $S_1 \times \cdots \times S_n$ of simple factors, but it need not be closed in G.

Theorem 2 ([8])**.** *Let G be a connected Lie group, let $G = RS$ denote a Levi decomposition, and suppose that S is locally isomorphic to the product $S_1 \times \cdots \times S_n$ of connected simple factors. Then the following are equivalent:*

(i) *the group G has the AP,*
(ii) *the group S has the AP,*
(iii) *the groups S_i, with $i = 1, \ldots, n$, have the AP,*
(iv) *the real rank of the groups S_i, with $i = 1, \ldots, n$, is at most 1.*

Proof. The equivalence of (iii) and (iv) is [7, Theorem 5.1].

Suppose (iv) holds. We show that (i) and (ii) hold. The proof of this implication is the proof from [8]. For completeness, we include it: The group R is solvable, so it has the AP. Since the AP is preserved under group extenstions [9, Theorem 1.15], it is therefore enough to prove that G/R has the AP. The Lie algebra of G/R is the Lie algebra of S, so to prove (i) and (ii), it suffices to prove that any connected Lie group S' locally isomorphic to $S_1 \times \cdots \times S_n$ has the AP. Using [9, Theorem 1.15] again, we may assume that the center of S' is trivial. If Z_i denotes the center of S_i, then $S' \simeq (S_1/Z_1) \times \cdots \times (S_n/Z_n)$. Each group S_i/Z_i has real rank at most 1 and hence has the AP [3, 10]. Hence S' has the AP.

Suppose (iv) does not hold. We show that (i) does not hold (which also proves that (ii) does not hold). Fix some i. The closure $\overline{S}_i$ of S_i in G is of the form S_iC for some connected, compact, central subgroup of G (see [13, p. 614]). Let $G' = G/C$, and let $\pi\colon G \to G'$ be the quotient homomorphism. As C is compact, π is a closed map, and $\pi(S_i) = \pi(\overline{S}_i)$ is a closed subgroup of G'. Since $\ker\pi \cap S_i$ is contained in the center of S_i and therefore is discrete (in the topology of S_i), the group $\pi(S_i)$ is locally isomorphic to S_i. If now the real rank of S_i is at least 2, then $\pi(S_i)$ does not have the AP [7, Theorem 5.1]. Hence G' does not have the AP, and it follows that G does not have the AP. □

A CHARACTERIZATION OF ALMOST CONNECTED GROUPS WITH THE AP

Let G be a locally compact group. Recall from [8] that there is a unique left invariant mean on $M_0A(G)$, and we say that G has property (T*) if this mean is weak* continuous. It is clear that non-compact groups with property (T*) do not have the AP. We establish the following converse for almost connected groups.

Theorem 3. *For an almost connected locally compact group G, the following are equivalent.*

(1) *G has the AP.*
(2) *No closed non-compact subgroup of G has property (T*).*

Proof. (1) $\Longrightarrow$ (2) is clear and holds also without the assumption of almost connectedness. We prove (2) $\Longrightarrow$ (1). Suppose G does not have AP. We prove the existence of a closed non-compact subgroup of G with property (T*).

As G is almost connected, there is a compact normal subgroup K of G such that G/K is a Lie group (see [12, Theorem 4.6]). Then G/K is a connected Lie group without the AP. It follows from the remark after Theorem C in [8] that we can find a closed, non-compact subgroup $H_0 \leq G/K$ with property (T*). Let H be the inverse image of H_0 in G. By [8, Proposition 5.13], H has property (T*), and H is clearly a closed non-compact subgroup of G. □

Remark 4. Theorem 3 is not true in general without the assumption of almost connectedness. In fact, there are discrete groups without the AP and without infinite subgroups with property (T*). This is not a new result, but simply a compilation of known results, the last ingredient being the recent result of Arzhantseva and Osajda (see [14, Theorem 2] and [1]) about the existence of discrete groups without property A but with the Haagerup property. Let G be such a group.

As property (T*) implies property (T) (see [8, Proposition 5.3]), and property (T) is an obstruction to the Haagerup property, it is clear that G has no infinite subgroups with property (T*).

On the other hand, if the discrete group G had the AP then its reduced group C*-algebra would have the strong operator approximation property. This is known to imply exactness of the group C*-algebra, which implies that G is an exact group. Finally, exactness for groups is equivalent to property A. This shows that G cannot have the AP. Proofs of all the claimed statements can be found in [2, Chapters 5 and 12].

REFERENCES

[1] Goulnara Arzhantseva and Damian Osajda. Graphical small cancellation groups with the Haagerup property. Preprint, arXiv:1404.6807, 2014.
[2] Nathanial P. Brown and Narutaka Ozawa. *C^*-algebras and finite-dimensional approximations*, volume 88 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2008.
[3] Michael Cowling and Uffe Haagerup. Completely bounded multipliers of the Fourier algebra of a simple Lie group of real rank one. *Invent. Math.*, 96(3):507–549, 1989.
[4] Jean de Cannière and Uffe Haagerup. Multipliers of the Fourier algebras of some simple Lie groups and their discrete subgroups. *Amer. J. Math.*, 107(2):455–500, 1985.
[5] Pierre Eymard. L'algèbre de Fourier d'un groupe localement compact. *Bull. Soc. Math. France*, 92:181–236, 1964.
[6] Uffe Haagerup and Tim de Laat. Simple Lie groups without the Approximation Property. *Duke Math. J.*, 162(5):925–964, 2013.

[7] Uffe Haagerup and Tim de Laat. Simple Lie groups without the Approximation Property II. *Trans. Amer. Math. Soc.*, 368(6):3777–3809, 2016.
[8] Uffe Haagerup, Søren Knudby, and Tim de Laat. A complete characterization of connected Lie groups with the Approximation Property. Preprint, to appear in Ann. Sci. École Norm. Sup., arXiv:1412.3033, 2014.
[9] Uffe Haagerup and Jon Kraus. Approximation properties for group C^*-algebras and group von Neumann algebras. *Trans. Amer. Math. Soc.*, 344(2):667–699, 1994.
[10] Mogens Lemvig Hansen. Weak amenability of the universal covering group of $\mathrm{SU}(1, n)$. *Math. Ann.*, 288(3):445–472, 1990.
[11] Vincent Lafforgue and Mikael De la Salle. Noncommutative L^p-spaces without the completely bounded approximation property. *Duke Math. J.*, 160(1):71–116, 2011.
[12] Deane Montgomery and Leo Zippin. *Topological transformation groups*. Interscience Publishers, New York-London, 1955.
[13] George Daniel Mostow. The extensibility of local Lie groups of transformations and groups on surfaces. *Ann. of Math. (2)*, 52:606–636, 1950.
[14] Damian Osajda. Small cancellation labellings of some infinite graphs and applications. Preprint, arXiv:1406.5015, 2014.
[15] V. S. Varadarajan. *Lie groups, Lie algebras, and their representations*, volume 102 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1984. Reprint of the 1974 edition.

SØREN KNUDBY
MATHEMATICAL INSTITUTE, UNIVERSITY OF MÜNSTER,
EINSTEINSTRASSE 62, 48149 MÜNSTER, GERMANY.

E-mail address: knudby@uni-muenster.de